

VII REFERÊNCIAS BIBLIOGRÁFICAS

[1] Semanário Expresso, Caderno de Economia, 19 de Agosto de 2006

[2] Heslop RB, Jones H. “Química Inorgânica”, editado por Fundação Calouste Gulbenkian, 2ªed., Lisboa, 1987

[3] National Research Council (NRC) . Toxicological effects of Methylmercury. National Academy Press, Washington, DC. 2000

[4] World Health Organization. Toxnet – Methylmercury. Env. Health Criteria. 1990; 101:144

[5] Scientific Advisory Committee on Nutrition (SACN) and Committee on Toxicity (COT). Advice on fish consumption: benefits & risks. 2004

[6] Clarkson TW. The three modern faces of mercury. Environ Health Perspect. 2002; 110 (1):11-23

[7] Secretaria de Saúde do Departamento e Grupo de Química Ambiental e Ocupacional da Universidade de Cartagena, Faculdade de Ciências Químicas e Farmacêuticas, Cartagena, Colombia. Mercurio: contaminante ambiental. 2005

[8] Nazario B, Boyles S. Exposure to Thimerosal in the Womb Does Not Raise Risk of Autism. WebMD Medical News. 2007

[9] U.S. Department of Health and Human Services and U.S. Environmental Protection Agency. What you need to know about mercury in fish and shellfish, 2004

[10] Jornal Oficial da União Europeia. Regulamento (CE) nº 1881/2006 da Comissão, de 19 de Dezembro de 2006, que fixa os teores máximos de certos contaminantes presentes nos géneros alimentícios. Comissão das Comunidades Europeias. 2006

[11] World Health Organization. UN Committee recommends new dietary intake limits for mercury. 2003

[12] U.S. Environmental Protection Agency. Integrated Risk information System. Methylmercury. 2001

[13] EFSA. Opinion Of The Scientific Panel On Contaminants In The Food Chain On A Request From The Commission Related To Mercury And Methylmercury In Food. The EFSA Journal 2004; 34:1 -14

[14] Crinnion WJ. Environmental Medicine, Part Three: Long-Term effects of chronic low-dose mercury exposure. Altern Med Rev. 2000; 5(3):209-223)

[15] Castoldi AF, Coccini T, Ceccatelli S, Manzo L. Neurotoxicity and molecular effects of methylmercury. *Brain Res Bull.* 2001; 55:197-203.

[16] Zahir F, Rizwi S, Haq S, Khan R. Low dose mercury toxicity and human health. *Environ. Toxicol. & Pharmacol.* 2005; 20:351-360

[17] Mahaffey KR. Fish and shellfish as dietary sources of methylmercury and the ω -3 fatty acids, eicosahexaenoic acid and docosahexaenoic acid: risks and benefits. *Environ Res.* 2004; 95:414-428

[18] Stern AH. A review of the studies of the cardiovascular health effects of methylmercury with consideration of their suitability for risk assessment. *Environ Res.* 2005; 98:133–142

[19] Guallar E, Sanz-Gallardo MI, van't Veer P, Bode P, Aro A, Gomez-Aracena J, Kark JD, Riemersma RA, Martin-Moreno JM, Kok FJ; Heavy Metals and Myocardial Infarction Study Group. Mercury, fish oils, and the risk of myocardial infarction. *N Engl J Med.* 2002; 347:1747-1754.

[20] EFSA. Opinion Of The Scientific Panel On Contaminants In The Food Chain On A Request From The European Parliament Related To The Safety Assessment Of Wild And Farmed Fish. *The EFSA Journal* 2005; 236:1 -118

[21] Sidhu KS. Health benefits and potential risks related to consumption of fish or fish oil. Regul Toxicol Pharmacol. 2003; 38:336-344

[22] Collette RL. Fish: Nutritional benefits and current issues. IFT Annual Meeting - Nutraceutical & Functional Foods Division, 2005

[23] Chapman L, Chan HM. The influence on methylmercury intoxication. Environ Health Perspect. 2000; 108(suppl1):29-56

[24] A.I. Cabañero, C.M.L. Carvalho, Y. Madrid, M.C.C. Batoréu and C. Cámara. "Quantification and Speciation of Mercury and Selenium in fish Samples of High Consumption in Spain and Portugal" Biol. Trace Elem. Res., 2005, 103, 17-35

[25] Santos APM, Mateus ML, Carvalho CML, Batoréu MCC. Biomarkers of exposure and effect as indicators of the interference of selenomethionine on methylmercury toxicity. Toxicol.Lett. 2007; 169: 121–128

[26] Mozaffarian D, Lemaitre RN, Kuller LH, Burke GL, Tracy RP, Siscovick DS. Cardiac Benefits of Fish Consumption May Depend on the Type of Fish Meal Consumed. Circulation 2003; 107:2632-s34

[27] Schmidt E, Arnesen H, Caterina R, Rasmussen L, Kristensen S. Marine n-3 polyunsaturated fatty acids and coronary heart disease Part I. Background,

epidemiology, animal data, effects on risk factors and safety. *Thrombosis Research* 2005; 115:163-170

[28] Marszalek JR, and Lodish HF. Docosahexaenoic Acid, Fatty Acid–Interacting Proteins, and Neuronal Function: Breastmilk and Fish Are Good for You. *Annu. Rev. Cell Dev. Biol.* 2005; 21:633-57

[29] Agostoni C, Verduci E. DHA in pregnancy benefits child development. *Pediatr Res.* 2003;54:292-293

[30] Oh R. Practical Applications of Fish Oil (Omega-3 fatty acids) in Primary Care. *J Am Board Fam Pract.* 2005; 18:28-36.

[31] Bakker EC, Ghys AJ, Kester AD, Vles JS, Dubas JS, Blanco CE, Hornstra G. Long-chain polyunsaturated fatty acids at birth and cognitive function at 7 y of age. *Eur J Clin Nutr.* 2003; 57:89-95.

[32] Ghys A, Bakker E, Hornstra G, van den Hout M. Red blood cell and plasma phospholipid arachidonic and docosahexaenoic acid levels at birth and cognitive development at 4 years of age. *Early Hum Dev.* 2002; 69:83-90.

[33] Olsen SF, Secher NJ. Low consumption of seafood in early pregnancy as a risk factor for preterm delivery: prospective cohort study. *BMJ.* 2002; 324:447.

[34] American Heart Association Scientific Statement. Fish, levels of mercury and omega-3 fatty acids. 2006

[35] Simopoulos A. Omega-3 Fatty Acids in Inflammation and Autoimmune Diseases. *J Am Coll Nutr.* 2002; 21(6):495-505

[36] Solfrizzi V, D'Introno A, Colacicco AM, Capurso C, Del Parigi A, Capurso S, Gadaleta A, Capurso A, Panza F. Dietary fatty acids intake: possible role in cognitive decline and dementia. *Exp Gerontol.* 2005; 40:257–270

[37] European Commission – DG Environment. Position paper on mercury: The technical report of the European air quality directive on mercury. 2003

[38] U.S. Department of Health and Human Services and U.S. Environmental Protection Agency. Mercury Levels in Commercial Fish and Shellfish, 2004.

[39] C.M.L. Carvalho, A.I.N.M. Matos, M.L. Mateus, A.P.M. Santos and M.C.C. Batoréu. "High-Fish Consumption and Risk Prevention: Exposure Assessment to Methylmercury in Portugal" (submitted)

[40] Björnberg KA, Vahter M, Grawé KP, Berglund M. Methyl mercury exposure in Swedish women with high fish consumption. *Sci Total Environ.* 2005; 341:45-52

[41] National Food Administration. Good advice for pregnant or breast-feeding women.

2004 disponível em: http://www.slv.se/templates/SLV_page.aspx?id=7035